

On chromatic number of Latin square graphs

Nazli Besharati^a, Luis Goddyn^b, E.S. Mahmoodian^c, M. Mortezaeefar^c

^a*Department of Mathematical Sciences, Payame Noor University, P.O. Box 19395-3697, Tehran, I. R. Iran*

^b*Department of Mathematics, Simon Fraser University, Burnaby, BC V5A 1S6 Canada*

^c*Department of Mathematical Sciences, Sharif University of Technology, P. O. Box 11155-9415, Tehran, I. R. Iran*

Abstract

The chromatic number of a Latin square is the least number of partial transversals which cover its cells. This is just the chromatic number of its associated Latin square graph. Although Latin square graphs have been widely studied as strongly regular graphs, their chromatic numbers appear to be unexplored. We determine the chromatic number of a circulant Latin square, and find bounds for some other classes of Latin squares. With a computer, we find the chromatic number for all main classes of Latin squares of order at most eight.

Keywords: Transversal, Partial transversal, Latin square, Latin square graphs, Net graphs, Coloring, Cayley table, Orthogonal array, Orthogonal Mate.

2000 MSC: 05B15- 05C15

1. Introduction and preliminaries

Let \mathbf{L} be a Latin square of order n . The chromatic number of \mathbf{L} , denoted by $\chi(\mathbf{L})$, is the minimum number of partial transversals of \mathbf{L} which together cover the cells of \mathbf{L} . Since each partial transversal uses at most n of the n^2 cells in \mathbf{L} , we observe the following.

Proposition 1. *Every Latin square \mathbf{L} of order n satisfies $\chi(\mathbf{L}) \geq n$, with equality holding if and only if \mathbf{L} has an orthogonal mate.*

Therefore $\chi(\mathbf{L})$ serves as a measure of how close \mathbf{L} is to having an orthogonal mate. We are surprised that reference to this natural invariant seems to be

absent from the substantial literature regarding transversals and orthogonality of Latin squares¹. An early version of some of our results appears in the M. Sc. thesis [12, in Farsi, Persian language] of the fourth author, under the supervision of the third author. In this paper, we find some bounds on $\chi(\mathbf{L})$ for general Latin squares and special classes such as complete Latin squares, Cayley tables of groups, circulants, and all Latin squares of order at most eight.

For the definitions not given here one may refer to [1] and [4]. Let \mathbf{L} be a Latin square of order n with cells $\{(r, c) \mid r, c \in \{0, 1, 2, \dots, n-1\}\}$; each cell contains a **symbol** from an alphabet of size n , and no row or column of \mathbf{L} contains a repeated symbol. A cell (r, c) containing the symbol $s = \mathbf{L}_{r,c}$ is sometimes represented by the triple (r, c, s) . A **partial transversal of length k** is a set of k cells, where no two cells have the same row, column or symbol. A **transversal** is a partial transversal of length n . The Latin square \mathbf{L} has an **orthogonal mate** if and only if it has a decomposition into disjoint transversals. We say that \mathbf{L} is **row-complete** if every ordered pair of distinct symbols appears (exactly once) in the set

$$\{(s, s') \mid (r, c, s), (r, c+1, s') \in \mathbf{L}, \text{ for } 0 \leq r \leq n-1 \text{ and } 0 \leq c \leq n-2\}.$$

The **Latin square graph** of \mathbf{L} is the simple graph $\Gamma(\mathbf{L})$ whose vertices are the cells of \mathbf{L} , and where distinct cells (r, c, s) and (r', c', s') are adjacent if (exactly) one of the equations $r = r'$, $c = c'$, $s = s'$ is satisfied. Accordingly, each edge of $\Gamma(\mathbf{L})$ is called, respectively, a **row edge**, a **column edge** or a **symbol edge**. Latin square graphs were introduced by R. C. Bose [2] as examples of strongly regular graphs; see [8, Section 10.4] for further discussion. Bose used the notation $L_3(n)$ for this graph. But this notation does not specify the Latin square from which the graph arises. So we use the notation $\Gamma(\mathbf{L})$ for the graph corresponding to the given Latin square \mathbf{L} .

The independent sets of $\Gamma(\mathbf{L})$ are the partial transversals of \mathbf{L} , and $\chi(\mathbf{L})$ is the chromatic number of $\Gamma(\mathbf{L})$. The isomorphism class of $\Gamma(\mathbf{L})$ is not affected by relabelling the rows, columns or symbols of \mathbf{L} , nor is it changed by applying a fixed permutation to the coordinates of every triple (r, c, s) in \mathbf{L} . Thus $\chi(\mathbf{L})$ is an invariant of the **main class** of \mathbf{L} .

Let (G, \circ) be a finite group of order n . A **Cayley table** for G is an $n \times n$ matrix, denoted L_G , where the cell (i, j) contains the group element $g_i \circ g_j$,

¹See the Addendum for further details

for some fixed enumeration $G = \{g_0, \dots, g_{n-1}\}$. It is easy to see that L_G is a Latin square. If G is a cyclic group, then L_G is called a **circulant Latin square**. Figure 1 shows the graph of the circulant $L_{\mathbb{Z}_3}$.




Figure 1: The graph $\Gamma(L_{\mathbb{Z}_3})$ with each vertex (r, c, s) labelled by s .

We summarize the results of this paper. Let \mathbf{L} be a Latin square of order n .

- $n \leq \chi(\mathbf{L}) \leq 3n - 2$.
- If \mathbf{L} is row-complete, then $\chi(\mathbf{L}) \leq 2n$.
- For large n we have $\chi(\mathbf{L}) = n + o(n)$.
- For every group G of order n , either $\chi(L_G) = n$ or $\chi(L_G) \geq n + 2$.
- If \mathbf{L} is a circulant (that is $\mathbf{L} \cong L_{\mathbb{Z}_n}$), then $\chi(\mathbf{L}) = \begin{cases} n & \text{if } n \text{ is odd} \\ n + 2 & \text{if } n \text{ is even.} \end{cases}$
- If $n \leq 8$, then

$$\chi(\mathbf{L}) \leq \begin{cases} n + 1 & \text{if } n \text{ is odd} \\ n + 2 & \text{if } n \text{ is even.} \end{cases} \quad (1)$$

We propose the following.

Conjecture 1. *Every Latin square \mathbf{L} satisfies (1).*

Conjecture 1 would surely be challenging to prove, even for Cayley tables of groups. Since $(n + 1)(n - 1) < n^2$ and $(n + 2)(n - 2) < n^2$, every Latin square \mathbf{L} which satisfies (1) must have a transversal (if n is odd) or a partial transversal of length $n - 1$ (if n is even). So Conjecture 1 would imply two long-standing conjectures of Brualdi-Stein and Ryser.

Conjecture 2. ([5, 15]) *Every Latin square of even order n contains a partial transversal of length $n - 1$.*

Conjecture 3. ([14]) *Every Latin square of odd order contains a transversal.*

2. Some Upper Bounds

The graph of a Latin square \mathbf{L} of order n is regular of degree $3n - 3$. This immediately gives $\chi(\mathbf{L}) \leq 3n - 2$. We can improve this bound in case \mathbf{L} has additional structure.

A k -plex is a set of kn cells which has k representatives from each row and each column and each symbol of \mathbf{L} . A (k_1, k_2, \dots, k_d) -partition is a partition K_1, K_2, \dots, K_d where each K_i is a k_i -plex.

Proposition 2. *If a Latin square \mathbf{L} of order n has a (k_1, k_2, \dots, k_d) -partition, then $\chi(\mathbf{L}) \leq 3n - 2d$.*

Proof. Let K_1, K_2, \dots, K_d be a (k_1, k_2, \dots, k_d) -partition of \mathbf{L} . For each i , the induced subgraph $\Gamma(\mathbf{L})[K_i]$ is regular of degree $3k_i - 3$, so it is $(3k_i - 2)$ -colorable. Thus, $\chi(\mathbf{L}) \leq \sum_{i=1}^d (3k_i - 2) = 3n - 2d$. ■

Wanless [16, 17] has conjectured that every Latin square of order n has a (k_1, k_2, \dots, k_d) -partition where $d \geq \lfloor \frac{n}{2} \rfloor$. If true, his conjecture would improve the general upper bound to $\chi(\mathbf{L}) \leq 2n + 1$.

Corollary 1. *Let t be the maximum number of disjoint transversals in a Latin square \mathbf{L} of order n . Then*

$$\begin{cases} \chi(\mathbf{L}) \leq 3n - 2t - 2 & \text{if } t \leq n - 2 \\ \chi(\mathbf{L}) = n & \text{otherwise.} \end{cases}$$

A Latin square \mathbf{L} of order n is **row-complete** (sometimes it is called a Roman square) if the ordered pairs $(\mathbf{L}_{i,j}, \mathbf{L}_{i,j+1})$ are all distinct for $0 \leq i \leq n - 1$ and $0 \leq j \leq n - 2$. Row-complete Latin squares are used in the design of sequential experiments [5].

Proposition 3. *If \mathbf{L} is a row-complete Latin square of order n , then $\chi(\mathbf{L}) \leq 2n$.*

0 ₁	1 ₃	3 ₂	2 ₄
1 ₂	2 ₀	0 ₃	3 ₅
2 ₃	3 ₁	1 ₀	0 ₆
3 ₀	0 ₂	2 ₁	1 ₇

Figure 2: Subscripts indicate an 8-coloring of row-complete Latin square of order 4.

Proof. We color each cell $\mathbf{L}_{i,j}$, $0 \leq i \leq n-1$ and $0 \leq j \leq n-2$, with the entry of $\mathbf{L}_{i,j+1}$. By the definition of row-complete, the cells with color c are a partial transversal in \mathbf{L} . We color the last column of \mathbf{L} with n new colors. This gives $\chi(\mathbf{L}) \leq 2n$. See Figure 2 for an example. ■

The above bounds are far from optimal for large Latin squares. Let V be the disjoint union $R \cup C \cup S$ where R, C, S is the set of rows, columns, and symbols of a Latin square \mathbf{L} of order n . Let $\mathcal{H} = (V, E)$ be the 3-uniform 3-partite hypergraph, where there is a hyperedge $\{r, c, s\} \in E$ for every cell $(r, c, s) \in \mathbf{L}$. Then \mathcal{H} is a linear hypergraph i.e., no two hyperedges share more than one vertex, and \mathcal{H} is regular of degree n . Also $\chi(\mathbf{L})$ is the chromatic index of \mathcal{H} , the least number of colors needed to color E so that no two adjacent hyperedges get the same color. The following general result of Molloy and Reed [11] implies an asymptotically optimal bound for $\chi(\mathbf{L})$.

Theorem 1. *For every k there exists a constant c_k such that every k -uniform linear hypergraph of maximum degree n has chromatic index at most $n + c_k(\log n)^4 n^{1-1/k}$.*

Corollary 2. *As $n \rightarrow \infty$ every Latin square \mathbf{L} of order n satisfies $\chi(\mathbf{L}) \leq n + o(n)$.*

3. Chromatic Number of Cayley Tables

The set of finite groups G for which $\chi(L_G) = |G|$ has been recently characterized.

Theorem 2. *For any finite group G of order n with identity element ϵ , the following are equivalent.*

1. $\chi(L_G) = n$
2. $\chi(L_G) \leq n + 1$.

3. L_G has a transversal.
4. For some enumeration g_1, g_2, \dots, g_n of G we have $g_1 g_2 \dots g_n = \epsilon$.
5. Every Sylow 2-subgroup of G is either trivial or non-cyclic.

Proof. The equivalence of statements 1. and 3. is classic, while the equivalence of 4. and 5. was shown in [6]. The equivalence of 3. and 4. was conjectured by Hall and Paige [9], and proved by Bray, Evans and Wilcox using the classification of finite simple groups (see [7]). Trivially, 1. implies 2. Also, 2. implies 3., as per the discussion before Conjecture 2 ■.

Corollary 3. *We have*

1. $\chi(L_G) = |G|$ for every group G of odd order (also see [3]).
2. For every group G of order n , either $\chi(L_G) = n$ or $\chi(L_G) \geq n + 2$.
3. Let G be an Abelian group of order n , $\chi(L_G) \geq n + 2$ if and only if G has a unique element of order 2 (also see [13]).

The rest of this section is devoted to proving that the lower bound of Corollary 3(3) is tight for cyclic groups.

Theorem 3. *For $n \geq 1$ the circulant Latin square of order n satisfies*

$$\chi(L_{\mathbb{Z}_n}) = \begin{cases} n & \text{if } n \text{ is odd} \\ n + 2 & \text{if } n \text{ is even.} \end{cases}$$

Our proof of Theorem 3 utilizes the following class of graphs.

Definition. *The Möbius ladder of order $2n$ is the cubic graph M obtained from a cycle C of length $2n$ by adding n new edges, each connecting an opposite pair of vertices of C . The cycle C is called the **rim** of M , and the added edges are called the **rungs** of M . (See Figure 3 and Figure 4(b)). Two vertices of M are said to be **nearly antipodal** in M if they are at distance $n - 1$ in C .*

The following proposition is straightforward to verify.

Proposition 4. *If x, x' are nearly antipodal vertices of a Möbius ladder M , then $M - \{x, x'\}$ is bipartite.*




Figure 3: A Möbius ladder of order 16.

Let $L_{\mathbb{Z}_n} = \{(r, c, s) \in \mathbb{Z}_n \times \mathbb{Z}_n \times \mathbb{Z}_n \mid s = r + c \pmod{n}\}$ be the circulant Latin square of order n . For $0 \leq i < n$, we define the i th right diagonal of $L_{\mathbb{Z}_n}$ to be the set $T_i = \{(r, r + i, 2r + i) \mid 0 \leq r \leq n - 1\}$.

Proposition 5. *For $0 \leq i < n$, the subgraph $\langle T_i \cup T_{i+1} \rangle$ of $\Gamma(L_{\mathbb{Z}_n})$ induced by the 2-plex $T_i \cup T_{i+1}$ is isomorphic to the Möbius ladder of order $2n$. Moreover if $n = 2m$ is even, then every cell $x \in T_i$ is nearly antipodal to each of the cells $x' = x + (m, m + 1, 1)$ and $x'' = x + (m - 1, m, -1)$.*

Proof. The row edges and column edges in $\langle T_i \cup T_{i+1} \rangle$ comprise the rim of the Möbius ladder. For each $(r, c, s) \in T_i$, we have $c - r = i$, $c + r = s$, and the only other solution to the system $\{c' - r' \in \{i, i + 1\}, c' + r' = s\} \pmod{n}$ is $(r', c') = (r + \lfloor \frac{n}{2} \rfloor, c + \lfloor \frac{n}{2} \rfloor)$. Therefore each symbol edge in $\langle T_i \cup T_{i+1} \rangle$ is a rung joining an opposite pair of vertices along the rim of $\langle T_i \cup T_{i+1} \rangle$. If $n = 2m$ and $x \in T_i$, then both x' and x'' are nearly antipodal to x since they are adjacent to $x + (m, m, 0)$ along the rim of $\langle T_i \cup T_{i+1} \rangle$, and $\{x, x + (m, m, 0)\}$ is a rung of $\langle T_i \cup T_{i+1} \rangle$. ■

See Figure 4 for an example.


Proof of Theorem 3. If n is odd it follows by Corollary 3(1).

If n is even, by Corollary 3(3), it suffices to show that $\Gamma(L_{\mathbb{Z}_{2m}})$ has a proper $(2m + 2)$ -coloring for $m \geq 1$. This is true for $m = 1$ since $\Gamma(L_{\mathbb{Z}_2}) \cong K_4$. We assume $m \geq 2$ and let $k = \lceil \frac{m}{2} \rceil$. Using the notation $[t] = \{0, 1, \dots, t - 1\}$, we define four sets of cells:

$$\begin{aligned} X &= \{x_j \mid j \in [k]\}, & X' &= \{x'_j \mid j \in [k]\}, \\ Y &= \{y_j \mid j \in [m - k]\}, & Y' &= \{y'_j \mid j \in [m - k]\}, \end{aligned}$$

0	1	2	3	<u>4</u>	5	6	7
1	2	3	4	5	6	7	0
2	3	4	5	6	7	0	1
<u>3</u>	4	5	6	7	0	1	2
4	5	6	7	0	1	2	3
5	6	7	0	1	2	3	4
6	7	0	1	2	3	4	5
7	0	1	2	3	4	5	6

(a)



(b)

Figure 4: For $L_{\mathbb{Z}_8}$ we indicate $T_4 \cup T_5$ in bold face, and underline the near antipodal cells $x = (0, 4, 4)$ and $x'' = x + (3, 4, -1) = (3, 0, 3)$ in the Möbius ladder $\langle T_4 \cup T_5 \rangle$.

where

$$\begin{aligned} x_j &= (j, 3j, 4j), & x'_j &= x_j + (d_r, d_c, d_s), & \text{for } j \in [k], \\ y_j &= x_j + (0, 2k, 2k), & y'_j &= y_j + (d_r, d_c, d_s), & \text{for } j \in [m - k], \end{aligned}$$

and where

$$(d_r, d_c, d_s) = \begin{cases} (m, m + 1, 1) & \text{if } m \equiv 0 \pmod{3}, \\ (m - 1, m, -1) & \text{if } m \not\equiv 0 \pmod{3}. \end{cases}$$

Each of these sets is (essentially) a translation of X :

$$\begin{aligned} X' &= X + (d_r, d_c, d_s), & (2) \\ Y \cup Y' &= \begin{cases} (X \cup X') + (0, 2k, 2k) & \text{if } m \text{ is even,} \\ (X \cup X' \setminus \{x_k, x'_k\}) + (0, 2k, 2k) & \text{if } m \text{ is odd.} \end{cases} & (3) \end{aligned}$$

See Figure 5 for examples.

Claim: Each of $X \cup X'$ and $Y \cup Y'$ is an independent set in the graph $\Gamma(L_{\mathbb{Z}_{2m}})$.

Proof of Claim: By (3) it suffices to prove the statement for $X \cup X' = \{x_j, x'_j \mid j \in [k]\}$. For any integer t , we define $\underline{t} \in [2m]$ to be the least positive residue of t modulo $2m$. For $j \in [k]$ we write $x_j = (r_j, c_j, s_j)$ and

0	1	2	3	4	5	6	7	8	9	10	11
1	2	3	4	5	6	7	8	9	10	11	0
2	3	4	5	6	7	8	9	10	11	0	1
3	4	5	6	7	8	9	10	11	0	1	2
4	5	6	7	8	9	10	11	0	1	2	3
5	6	7	8	9	10	11	0	1	2	3	4
6	7	8	9	10	11	0	1	2	3	4	5
7	8	9	10	11	0	1	2	3	4	5	6
8	9	10	11	0	1	2	3	4	5	6	7
9	10	11	0	1	2	3	4	5	6	7	8
10	11	0	1	2	3	4	5	6	7	8	9
11	0	1	2	3	4	5	6	7	8	9	10

0	1	2	3	4	5	6	7	8	9	10	11	12	13
1	2	3	4	5	6	7	8	9	10	11	12	13	0
2	3	4	5	6	7	8	9	10	11	12	13	0	1
3	4	5	6	7	8	9	10	11	12	13	0	1	2
4	5	6	7	8	9	10	11	12	13	0	1	2	3
5	6	7	8	9	10	11	12	13	0	1	2	3	4
6	7	8	9	10	11	12	13	0	1	2	3	4	5
7	8	9	10	11	12	13	0	1	2	3	4	5	6
8	9	10	11	12	13	0	1	2	3	4	5	6	7
9	10	11	12	13	0	1	2	3	4	5	6	7	8
10	11	12	13	0	1	2	3	4	5	6	7	8	9
11	12	13	0	1	2	3	4	5	6	7	8	9	10
12	13	0	1	2	3	4	5	6	7	8	9	10	11
13	0	1	2	3	4	5	6	7	8	9	10	11	12

Figure 5: The cells $X \cup X'$ (bold face) and $Y \cup Y'$ (underlined bold face) are shown for $L_{\mathbb{Z}_{12}}$ and for $L_{\mathbb{Z}_{14}}$.

$x'_j = (r'_j, c'_j, s'_j)$, and define the following six multisets:

$$\begin{aligned}
R &= \{\underline{r}_0, \underline{r}_1, \dots, \underline{r}_{k-1}\}, & R' &= \{\underline{r}'_0, \underline{r}'_1, \dots, \underline{r}'_{k-1}\}, \\
C &= \{\underline{c}_0, \underline{c}_1, \dots, \underline{c}_{k-1}\}, & C' &= \{\underline{c}'_0, \underline{c}'_1, \dots, \underline{c}'_{k-1}\}, \\
S &= \{\underline{s}_0, \underline{s}_1, \dots, \underline{s}_{k-1}\}, & S' &= \{\underline{s}'_0, \underline{s}'_1, \dots, \underline{s}'_{k-1}\}.
\end{aligned}$$

We aim to show that none of $R \cup R'$, $C \cup C'$, $S \cup S'$ has a repeated entry. For $j \in [k]$ each of the numbers $r_j = j$, $c_j = 3j$, $s_j = 4j$ is bounded above by $4(k-1) < 2m$, so we have

$$\underline{r}_j = r_j = j, \quad \underline{c}_j = c_j = 3j, \quad \underline{s}_j = s_j = 4j.$$

Therefore none of the lists R , C , S has a repeated entry. By (2), none of the lists R' , C' , S' has a repeated entry, so it suffices to show that each of the sets $R \cap R'$, $C \cap C'$, $S \cap S'$ is empty. First, since $m \geq 2$ we have $k < m$ so $R \subseteq \{0, 1, \dots, m-2\}$. For $j \in [k]$ we have $r'_j = r_j + d_r \in \{r_j + m, r_j + (m-1)\} \subseteq \{m-1, m, \dots, 2m-1\}$. Therefore $\underline{r}'_j = r'_j$ and $R \cap R' = \emptyset$. Second, for $j \in [k]$ we have $c'_j = c_j + d_c \in \{c_j + (m+1), c_j + m\}$. Therefore $0 \leq c'_j < 4m$ and

$$\underline{c}'_j = \begin{cases} 3j + d_c, & \text{if } c_j + d_c < 2m \\ 3j + d_c - 2m, & \text{if } c_j + d_c \geq 2m. \end{cases}$$

If $m \equiv 0 \pmod{3}$, then $d_c = m + 1$ so $\underline{c}'_j \equiv 1 \pmod{3}$. If $m \not\equiv 0 \pmod{3}$, then $d_c = m$ so $\underline{c}'_j \equiv \pm m \pmod{3}$. In both cases we find $\underline{c}'_j \not\equiv 0 \equiv \underline{c}_j \pmod{3}$ for $j \in [k]$. Therefore $C \cap C' = \emptyset$. Third, for $j \in [k]$ we have $s'_j = s_j \pm 1$ so s_j is even and s'_j is odd. Since $2m$ is even, we conclude that \underline{s}_j is even and \underline{s}'_j is odd for every $j \in [k]$. Therefore $S \cap S' = \emptyset$ and the claim is proved.

For every $j \in [m]$ we apply Proposition 5 to the 2-plex

$$M_j := T_{2j} \cup T_{2j+1},$$

where T_i is the i th right diagonal of $L_{\mathbb{Z}_{2m}}$. Each induced subgraph $\langle M_j \rangle$ is a Möbius ladder. For $j \in [k]$ we have that $x_j \in T_{2j}$, so the cell $x'_j = x_j + (d_r, d_c, d_s)$ is nearly antipodal to x_j in $\langle M_j \rangle$. Similarly, for $j \in [m - k]$ we find that y_j and y'_j are nearly antipodal vertices of $\langle M_{j+k} \rangle$. We now consider the following partition of $L_{\mathbb{Z}_{2m}}$ into $m + 1$ parts.

$$\{M_j - \{x_j, x'_j\} \mid j \in [k]\} \cup \{M_{j+k} - \{y_j, y'_j\} \mid j \in [m - k]\} \cup \{X \cup X' \cup Y \cup Y'\}$$

By Proposition 4 and the above claim, each part induces a bipartite subgraph of $\Gamma(L_{\mathbb{Z}_{2m}})$. We conclude that $\chi(L_{\mathbb{Z}_{2m}}) \leq 2m + 2$. ■

See Figure 6 for some examples.

0 ₄	1 ₁	2 ₅	3 ₃
1 ₅	2 ₀	3 ₄	0 ₂
2 ₃	3 ₂	0 ₁	1 ₀
3 ₀	0 ₃	1 ₂	2 ₁

0 ₆	1 ₁	2 ₃	3 ₂	4 ₇	5 ₅
1 ₅	2 ₀	3 ₁	4 ₆	5 ₃	0 ₄
2 ₄	3 ₅	4 ₀	5 ₁	0 ₂	1 ₃
3 ₃	4 ₄	5 ₇	0 ₀	1 ₆	2 ₂
4 ₂	5 ₆	0 ₅	1 ₄	2 ₁	3 ₀
5 ₀	0 ₃	1 ₂	2 ₅	3 ₄	4 ₁

0 ₈	1 ₁	2 ₂	3 ₃	4 ₉	5 ₅	6 ₆	7 ₇
1 ₆	2 ₀	3 ₁	4 ₈	5 ₂	6 ₄	7 ₅	0 ₉
2 ₇	3 ₆	4 ₀	5 ₁	6 ₃	7 ₂	0 ₄	1 ₅
3 ₉	4 ₇	5 ₆	6 ₀	7 ₈	0 ₃	1 ₂	2 ₄
4 ₅	5 ₄	6 ₇	7 ₉	0 ₁	1 ₀	2 ₃	3 ₈
5 ₃	6 ₅	7 ₄	0 ₆	1 ₇	2 ₁	3 ₀	4 ₂
6 ₂	7 ₃	0 ₅	1 ₄	2 ₆	3 ₇	4 ₁	5 ₀
7 ₀	0 ₂	1 ₃	2 ₅	3 ₄	4 ₆	5 ₇	6 ₁

Figure 6: Colorings of $L_{\mathbb{Z}_4}$, $L_{\mathbb{Z}_6}$, and $L_{\mathbb{Z}_8}$ by the method of Theorem 3. The colors are given as subscripts.

4. Latin squares of small orders

We have computed $\chi(\mathbf{L})$ for all main classes of order n , $2 \leq n \leq 7$. N. Shajari has verified by computer that all 283657 main classes of Latin squares of order 8 as listed in [10] are 10-colorable. The results are summarized in Table 1.

n	Number of main classes	Main class number as listed in [4]	χ
2	1	2.1 ($L_{\mathbb{Z}_2}$)	4
3	1	3.1 ($L_{\mathbb{Z}_3}$)	3
4	2	4.2 ($L_{\mathbb{Z}_2 \times \mathbb{Z}_2}$)	4
		4.1 ($L_{\mathbb{Z}_4}$)	6
5	2	5.1 ($L_{\mathbb{Z}_5}$)	5
		5.2	6
6	12	6.2, 6.3, 6.4, 6.5, 6.10, 6.11	7
		6.1, 6.6, 6.7, 6.8, 6.9, 6.12	8
7	147	7.3, 7.6, 7.7, 7.71, 7.105, 7.137	7
		All other main classes of order 7	8
8	283657	All main classes as listed in [10]	≤ 10

Table 1: Chromatic numbers of Latin squares of order n , $2 \leq n \leq 8$.

Acknowledgements. The work of Luis Goddyn was supported by NSERC Canada, while E.S. Mahmoodian was visiting Simon Fraser University and was partially supported by INSF.

Addendum. One of the referees for this paper asserts that $\chi(\mathbf{L})$ has been studied before by at least four different groups or individuals, he/she has discussed the invariant with some of these groups, and has seen it mentioned in several conference talks, but knows of no written work on the topic. Another referee informs us that some of our results are contained in a paper by Nicholas Cavenagh and Jaromy Kuhl that is the same concept but they call it “chromatic index” rather than “chromatic number”, and is about to be published in *Contributions to Discrete Mathematics* 12 (2016), issue 2.

References

In the references, each reference is followed by a number which is the page on which the reference is cited.

- [1] J. A. Bondy and U. S. R. Murty. *Graph theory*, volume 244 of *Graduate Texts in Mathematics*. Springer, New York, 2008. 2

- [2] R. C. Bose. Strongly regular graphs, partial geometries and partially balanced designs. *Pacific J. Math.*, 13:389–419, 1963. [2](#)
- [3] Richard A. Brualdi and Herbert J. Ryser. *Combinatorial matrix theory*, volume 39 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1991. [6](#), [13](#)
- [4] Charles J. Colbourn and Jeffrey H. Dinitz, editors. *Handbook of combinatorial designs*. Discrete Mathematics and its Applications (Boca Raton). Chapman & Hall/CRC, Boca Raton, FL, Second edition, 2007. [2](#), [11](#)
- [5] J. Dénes and A. D. Keedwell. *Latin squares and their applications*. Academic Press, New York, 1974. [4](#)
- [6] J. Dénes and A. D. Keedwell. A new conjecture concerning admissibility of groups. *European J. Combin.*, 10(2):171–174, 1989. [6](#)
- [7] Anthony B. Evans. The admissibility of sporadic simple groups. *J. Algebra*, 321(1):105–116, 2009. [6](#)
- [8] Chris Godsil and Gordon Royle. *Algebraic graph theory*, volume 207 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2001. [2](#)
- [9] Marshall Hall and L. J. Paige. Complete mappings of finite groups. *Pacific J. Math.*, 5:541–549, 1955. [6](#)
- [10] Brendan D. McKay, Alison Meynert, and Wendy Myrvold. Small Latin squares, quasigroups, and loops. *J. Combin. Des.*, 15(2):98–119, 2007. [10](#), [11](#)
- [11] Michael Molloy and Bruce Reed. Near-optimal list colorings. In *Proceedings of the Ninth International Conference “Random Structures and Algorithms” (Poznan, 1999)*, volume 17, pages 376–402, 2000. [5](#)
- [12] M. Mortezaeefar. Colorings of Latin Square Graphs and Block Designs. Master’s thesis, Sharif University of Technology, Tehran, Iran, 2009. (in Farsi, Persian language), <http://sdr.library.sharif.ir/resource/290994/-/&from=search&&query=-setfield=subjectkeyword&count=20&execute=true>. [2](#)

- [13] L. J. Paige. A note on finite Abelian groups. *Bull. Amer. Math. Soc.*, 53:590–593, 1947. [6](#)
- [14] H. J. Ryser. Neuere Probleme in der Kombinatorik (prepared by D.W. Miller). *Vortrage uber Kombinatorik*, pages 69–91, 1967. (cited in [\[3\]](#)). [4](#)
- [15] S. K. Stein. Transversals of Latin squares and their generalizations. *Pacific J. Math.*, 59(2):567–575, 1975. [4](#)
- [16] Ian M. Wanless. A generalisation of transversals for Latin squares. *Electron. J. Combin.*, 9(1):Research Paper 12, 15 pp. (electronic), 2002. [4](#)
- [17] Ian M. Wanless. Transversals in Latin squares: a survey. In *Surveys in combinatorics 2011*, volume 392 of *London Math. Soc. Lecture Note Ser.*, pages 403–437. Cambridge Univ. Press, Cambridge, 2011. [4](#)

nbesharati@pnu.ac.ir, goddyn@sfu.ca, emahmood@sharif.edu,
m.mortezaeefar@gmail.com